arXiv:0906.3037v1 [math.PR] 16 Jun 2009

On the density of the sum of two independent Student t-random vectors

C. Berg[a], C. Vignat[*,b]

[a] *Department of Mathematics, University of Copenhagen, Universitetsparken 5, DK-2100, Copenhagen, Denmark*
[b] *Institut Gaspard Monge, Université de Marne la Vallée, France*

Abstract

In this paper, we find an expression for the density of the sum of two independent $d-$dimensional Student $t-$random vectors $\mathbf{X}$ and $\mathbf{Y}$ with arbitrary degrees of freedom. As a byproduct we also obtain an expression for the density of the sum $\mathbf{N}+\mathbf{X}$, where $\mathbf{N}$ is normal and $\mathbf{X}$ is an independent Student $t-$vector. In both cases the density is given as an infinite series

$$\sum_{n=0}^{\infty} c_n f_n$$

where f_n is a sequence of probability densities on $\mathbb{R}^d$ and (c_n) is a sequence of positive numbers of sum 1, i.e. the distribution of a non-negative integer-valued random variable C, which turns out to be infinitely divisible for $d=1$ and $d=2$. When $d=1$ and the degrees of freedom of the Student variables are equal, we recover an old result of Ruben.

---

1. Introduction

The Student $t-$distributions plays an important role in statistics for the analysis of normal samples. One of the major properties of these distributions is their heavy-tail behaviour; the same behaviour is shown by the famous family of stable distributions. The Student t-distributions show the advantage of having an explicit expression, what allows for example to write down explicitly their likelihood function [4]; this is not the case for the stable distributions, except some well-known particular cases.

However, except in the Gaussian and Cauchy case, the Student t-distributions show the drawback of not being stable by convolution. Thus the study of their behavior by convolution is of high interest. This study has been first addressed in [15] and more recently in [17, 6], but concerns only the case where the degrees of freedom is an odd integer. The study of Student t-processes has recently given rise to a series of studies [8, 7, 11].

The $d-$dimensional Student $t-$density with $2\nu>0$ degrees of freedom is

$$f_\nu(\mathbf{t}) = A_{\nu,d}\left(1+\|\mathbf{t}\|^2\right)^{-\nu-\frac{d}{2}},\ \mathbf{t}\in\mathbb{R}^d;\ A_{\nu,d} = \frac{\Gamma\left(\nu+\frac{d}{2}\right)}{\Gamma(\nu)\,\pi^{\frac{d}{2}}}.$$

In the following, we provide an expression for the density of the sum

$$\mathbf{Y} = \mathbf{T}_1 + \mathbf{T}_2$$

where $\mathbf{T}_1$ and $\mathbf{T}_2$ are $d-$variate independent $t-$distributed random variables with respective degrees of freedom 2ν and 2μ. When $\mu=\nu$ and $d=1$, this equals an expression found by Ruben, cf. [14, formula 3.4]. As a byproduct, we also obtain an expression for the density of

$$\mathbf{Z} = \mathbf{X} + \mathbf{T},$$

---

[*] Corresponding author

Email addresses: `berg@math.ku.dk` (C. Berg), `vignat@univ-mlv.fr` (C. Vignat)

Preprint submitted to Annales de l'Institut Henri Poincaré (B) Probabilités et Statistiques *June 16, 2009*

where $\mathbf{X}$ is standard Gaussian. Moreover, we extend this result to the case of $\tilde{\mathbf{Z}} = a_1\mathbf{X} + a_2\mathbf{T}$ with $a_1, a_2 > 0$. In all cases, our expression is an infinite convex combination of a sequence of $d-$variate densities, which we describe next.

Let us introduce the family of densities

$$g_{k,\sigma}(\mathbf{z}) = \frac{1}{\sigma^d} g_{k,1}\left(\frac{\mathbf{z}}{\sigma}\right) = \frac{\Gamma(\frac{d}{2})}{\Gamma\left(k+\frac{d}{2}\right)\left(\sigma\sqrt{2\pi}\right)^d}\left(\frac{\|\mathbf{z}\|^2}{2\sigma^2}\right)^k \exp\left(-\frac{\|\mathbf{z}\|^2}{2\sigma^2}\right),\ \mathbf{z} \in \mathbb{R}^d \tag{1}$$

with $k \in \mathbb{N}$. We remark that $g_{0,\sigma}$ is the $d-$variate Gaussian density with zero mean and covariance matrix $\sigma^2 I_d$; for $k \neq 0$, by [9, Th. 2.9], $g_{k,\sigma}$ is the density of $\sigma\|\mathbf{X}_{2k+d}\|\mathbf{U}_d$ where $\mathbf{X}_{2k+d} \in \mathbb{R}^{2k+d}$ is a Gaussian vector with identity covariance matrix and $\mathbf{U}_d$ is uniformly distributed on the unit sphere $\mathcal{S}_d$ of $\mathbb{R}^d$ and independent of $\mathbf{X}_{2k+d}$.

Let us also introduce the family of densities

$$\phi_{n,\eta}\left(\mathbf{x}\right) = \frac{\Gamma\left(\frac{d}{2}\right)}{\pi^{\frac{d}{2}} B\left(\eta, n+\frac{d}{2}\right)} \frac{\|\mathbf{x}\|^{2n}}{\left(1+\|\mathbf{x}\|^2\right)^{\eta+n+\frac{d}{2}}},\ \mathbf{x} \in \mathbb{R}^d, \tag{2}$$

where $\eta > 0$. We note that, with the same notation as above, $\phi_{n,\eta}$ is the density of $\|\mathbf{T}_{2n+d}\|\mathbf{U}_d$, where $\mathbf{T}_{2n+d} \in \mathbb{R}^{2n+d}$ follows a multivariate Student t-distribution with 2η degrees of freedom

$$\frac{\Gamma\left(\eta+n+\frac{d}{2}\right)}{\Gamma\left(\eta\right)\pi^{n+\frac{d}{2}}}\left(1+\|\mathbf{x}\|^2\right)^{-\eta-n-\frac{d}{2}},\ \mathbf{x} \in \mathbb{R}^{2n+d}.$$

At last, we recall that a discrete random variable B follows a negative binomial distribution with parameters r and p (denoted as $NB\left(r,p\right)$) if

$$\Pr\left\{B = k\right\} = \binom{k+r-1}{k} p^r \left(1-p\right)^k,\ r > 0,\ 0 < p < 1,\ k = 0, 1 \ldots$$

NB distributions with non-integer parameter r are also known as Pólya distributions.

2. Results

The distribution of the sum of two independent Student $t-$vectors

$$\mathbf{Y} = \mathbf{T}_1 + \mathbf{T}_2$$

can be deduced by subordination from the distribution of $\mathbf{Z} = \mathbf{X} + \mathbf{T}$ where $\mathbf{X}$ is Gaussian : if $\tilde{\gamma}_\nu$ denotes the inverse Gamma measure with scale parameter $\frac{1}{4}$ and shape parameter $\nu > 0$

$$d\tilde{\gamma}_\nu\left(t\right) = \frac{1}{2^{2\nu}\Gamma\left(\nu\right)} \exp\left(-\frac{1}{4t}\right) t^{-\nu-1} dt,\ t > 0,$$

and if $g_t\left(\mathbf{x}\right)$ denotes the $d-$variate Gaussian semi-group of normal densities

$$g_t\left(\mathbf{x}\right) = \left(4\pi t\right)^{-\frac{d}{2}} \exp\left(-\frac{\|\mathbf{x}\|^2}{4t}\right),\ t > 0,\ \mathbf{x} \in \mathbb{R}^d,$$

then the Student $t-$density can be obtained as the mixture

$$f_\nu\left(\mathbf{x}\right) = \int_0^{+\infty} g_t\left(\mathbf{x}\right) d\tilde{\gamma}_\nu\left(t\right).$$

Therefore, we begin by finding an expression for the density of $\mathbf{Z}$.

2.1. The sum of a Gaussian and a Student t−vector

Our first result states as follows.

Theorem 1. *The density of* $\mathbf{Z} = \mathbf{X} + \mathbf{T}$ *can be written*

$$f_{\mathbf{Z}}(\mathbf{z}) = \sum_{k=0}^{+\infty} \alpha_k^{(\nu,\gamma)} g_{k,\sigma}(\mathbf{z}) \tag{3}$$

where the densities $g_{k,\sigma}$ *are defined in (1),* $\alpha_k^{(\nu,\gamma)}$ *are positive coefficients that sum to* 1 *and given by*

$$\alpha_k^{(\nu,\gamma)} = \frac{\Gamma(k+\frac{d}{2})}{k!\Gamma\left(\frac{d}{2}\right)\Gamma(\nu)} \gamma^{2k} \int_0^{+\infty} \exp(-a) a^{\nu+\frac{d}{2}-1} \left(a+\gamma^2\right)^{-k-\frac{d}{2}} da, \;\; k \geq 0 \tag{4}$$

and γ *is defined as*

$$\gamma = \frac{1}{\sigma\sqrt{2}}. \tag{5}$$

Proposition 2. *The same result as in Theorem 1 holds for* $\tilde{\mathbf{Z}} = a_1\mathbf{X} + a_2\mathbf{T}, \;\; a_1 \in \mathbb{R}^+, \, a_2 \in \mathbb{R}^+$ *replacing* σ *by* $a_1\sigma$ *and* γ *by* $\frac{a_2}{\sigma\sqrt{2}a_1}$.

Note that $a_1\mathbf{X}$ and $a_2\mathbf{T}$ have the same distributions as $(-a_1\mathbf{X})$ and $(-a_2\mathbf{T})$ respectively, so that we only need to consider the case $a_1 > 0$, $a_2 > 0$ in the above Proposition.

2.2. The sum of two Student t−vectors

Starting from the distribution (3) we obtain the following

Theorem 3. *The distribution of* $\mathbf{Y} = \mathbf{T}_1 + \mathbf{T}_2$, *where* $\mathbf{T}_1$ *and* $\mathbf{T}_2$ *are independent t−distributed with respective degrees of freedom* $2\nu > 0$ *and* $2\mu > 0$, *can be written as the infinite convex combination*

$$f_{\mathbf{Y}}(\mathbf{x}) = \sum_{n=0}^{+\infty} c_n^{(\nu,\mu)} \phi_{n,\nu+\mu}(\mathbf{x}), \tag{6}$$

where $\phi_{n,\nu+\mu}$ *are the probability densities given by (2) and* $c_n^{(\nu,\mu)}$ *are positive coefficients that sum to* 1 *defined by*

$$c_n^{(\nu,\mu)} = \frac{1}{B(\nu,\mu)} \frac{\left(\frac{d}{2}\right)_n}{n!} \int_0^1 t^{\mu+\frac{d}{2}-1} (1-t)^{\nu+\frac{d}{2}-1} \left(1-t+t^2\right)^n dt, \;\; n \geq 0. \tag{7}$$

Note that (7) shows that $c_n^{(\mu,\nu)} = c_n^{(\nu,\mu)}$ as is to be expected.

Remark 1. The Fourier transform of the Student t−density in the case $d = 1$ is

$$\hat{f}_\nu(y) = \int_{-\infty}^{+\infty} e^{ixy} f_\nu(x)\, dx = k_\nu(|y|) \tag{8}$$

with

$$k_\nu(u) = \frac{2^{1-\nu}}{\Gamma(\nu)} u^\nu K_\nu(u), \;\; u \geq 0,$$

where K_ν is the MacDonald function.

We claim that

$$D^{2n} \hat{f}_{\mu+\nu+n}(y) = (-1)^n \frac{\left(\frac{1}{2}\right)_n}{(\mu+\nu)_n} \hat{\phi}_{n,\mu+\nu}(y), \;\; y \in \mathbb{R}. \tag{9}$$

In fact, from (8), we get

$$D^{2n}\hat{f}_{\mu+\nu+n}\left(y\right)=\left(-1\right)^{n}\int_{-\infty}^{+\infty}e^{ixy}A_{\mu+\nu+n,1}\frac{x^{2n}}{\left(1+x^{2}\right)^{\mu+\nu+n+\frac{1}{2}}}dx$$

which is (9) because

$$A_{\mu+\nu+n,1}B\left(\mu+\nu,n+\frac{1}{2}\right)=\frac{\left(\frac{1}{2}\right)_{n}}{\left(\mu+\nu\right)_{n}}.$$

Theorem 3 can be expressed for $d=1$ as

$$k_{\nu}\left(|y|\right)k_{\mu}\left(|y|\right)=\sum_{n=0}^{+\infty}c_{n}^{(\nu,\mu)}\hat{\phi}_{n,\mu+\nu}\left(y\right)$$

hence by (9)

$$=\sum_{n=0}^{+\infty}(-1)^{n}c_{n}^{(\nu,\mu)}\frac{\left(\mu+\nu\right)_{n}}{\left(\frac{1}{2}\right)_{n}}k_{\mu+\nu+n}^{(2n)}\left(|y|\right)$$

and finally by (7), for $u>0$,

$$B\left(\mu,\nu\right)k_{\nu}\left(u\right)k_{\mu}\left(u\right)=\sum_{n=0}^{+\infty}\frac{\left(-1\right)^{n}\left(\mu+\nu\right)_{n}}{n!}C_{n}(\nu,\mu)k_{\mu+\nu+n}^{(2n)}\left(u\right)$$

with

$$C_{n}(\nu,\mu)=\int_{0}^{1}t^{\mu-\frac{1}{2}}\left(1-t\right)^{\nu-\frac{1}{2}}\left(1-t+t^{2}\right)^{n}dt.$$

In terms of the Macdonald function, this equation can be written

$$K_{\nu}\left(u\right)K_{\mu}\left(u\right)=\sum_{n=0}^{+\infty}\frac{\left(-1\right)^{n}}{n!2^{n+1}}C_{n}(\nu,\mu)u^{-\mu-\nu}D^{2n}\left\{u^{\mu+\nu+n}K_{\mu+\nu+n}\left(u\right)\right\}.$$

We have not been able to find this expression in the literature.
Still another interpretation of formula (6) for $d=1$ is to rewrite it as

$$f_{\mu}*f_{\nu}\left(x\right)=f_{\mu+\nu}\left(x\right)\sum_{n=0}^{+\infty}\frac{\left(\mu+\nu+\frac{1}{2}\right)_{n}}{\left(\frac{1}{2}\right)_{n}}c_{n}^{(\nu,\mu)}\left(\frac{x^{2}}{1+x^{2}}\right)^{n},$$

which shows that $\frac{f_{\mu}*f_{\nu}(x)}{f_{\mu+\nu}(x)}$ has a holomorphic extension to the open set $\left\{z\in\mathbb{C}\mid|\frac{z^{2}}{1+z^{2}}|<1\right\}$, which is the region between the two branches of the equilateral hyperbola $y^{2}-x^{2}=\frac{1}{2}$.

Remark 2. In [14], Ruben shows that in the $d=1$ dimensional case, the density f_{Z} of the convolution $Z=T_{1}+T_{2}$ can be written as a mixture of scaled Student t-distributions with $\mu+\nu$ degrees of freedom, i.e. as

$$f_{Z}(z)=\int_{2}^{+\infty}\frac{1}{t}f_{\mu+\nu}\left(\frac{z}{t}\right)h(t)dt$$

for some probability density h on $]2,+\infty[$. This representation differs from (6) which is a discrete mixture of the densities (2). In the same paper, Ruben obtains several expressions for the density $f_{\mathbf{Z}}$ in the case $\mu=\nu$. Ruben's formula (3.4) coincides with formula (6) in the special case $\mu=\nu$, $d=1$.

3. Properties of the mixing distributions

In this section, we derive several properties of the discrete distributions (4) and (7).

3.1. Stochastic interpretation

Since, as a consequence of (4), the coefficients $\alpha_k^{(\nu,\gamma)}$ are positive, and since they sum to 1 (see Theorem 7), a stochastic interpretation of Theorem 1 is as follows:

$$\mathbf{Z} = \mathbf{X} + \mathbf{T} \overset{d}{=} \mathbf{G}_K$$

where, given $K = k$, $\mathbf{G}_K$ has the conditional density (1) and K is a discrete random variable with

$$\Pr\{K = k\} = \alpha_k^{(\nu,\gamma)}, \;\; k = 0, 1, \ldots \tag{10}$$

We deduce the following

Proposition 4. *The sum* $\mathbf{Z} = \mathbf{X} + \mathbf{T}$ *is distributed as* $\sigma \|\mathbf{X}_{2K+d}\| \mathbf{U}_d$ *where*

- *vector* $\mathbf{U}_d$ *is uniformly distributed on the unit sphere* $\mathcal{S}_d$ *in* $\mathbb{R}^d$,
- *vector* $\mathbf{X}_{2k+d} \in \mathbb{R}^{2k+d}$ *is Gaussian with identity covariance matrix,*
- *random variable* K *defined by (10) follows a compound negative binomial distribution* $NB\left(\frac{d}{2}, \frac{a}{1+a}\right)$, *where the parameter* a *is Gamma distributed with scale parameter* γ^2 *and shape parameter* ν, *(in short* $a \sim \Gamma\left(\nu, \gamma^2\right)$*) so* a *has the density*
$$\frac{\gamma^{2\nu}}{\Gamma(\nu)} e^{-t\gamma^2} t^{\nu-1}, \; t > 0.$$

An analogous result in the case of the sum of two Student t-vectors is as follows: $\mathbf{Y} = \mathbf{T}_1 + \mathbf{T}_2$ is distributed as a random variable Φ_N such that, given $N = n$, Φ_N has the conditional density given by (2) and N is a discrete random variable defined by

$$\Pr\{N = n\} = c_n^{(\nu,\mu)}, \;\; n = 0, 1, \ldots \tag{11}$$

We deduce the following

Proposition 5. *The sum* $\mathbf{Y} = \mathbf{T}_1 + \mathbf{T}_2$ *is distributed as* $\|\mathbf{T}_{2N+d}\| \mathbf{U}_d$ *where*

- *vector* $\mathbf{U}_d$ *is uniformly distributed on the sphere* $\mathcal{S}_d$ *in* $\mathbb{R}^d$,
- *random vector* $\mathbf{T}_{2n+d} \in \mathbb{R}^{2n+d}$ *is Student t-distributed with* $2\nu + 2\mu$ *degrees of freedom*
- *random variable* N *defined by (11) follows a compound negative binomial distribution* $NB\left(\frac{d}{2}, t(1-t)\right)$, *where the parameter* t *follows a Beta distribution with parameters* μ *and* ν.

3.2. Hausdorff moment sequence

A sequence $\{s_n\}$ is called a Hausdorff moment sequence if it has the form

$$s_n = \int_0^1 x^n d\mu(x), \; n = 0, 1, \ldots$$

for some positive measure μ. It is called normalized if $\mu_0 = 1$. The following result holds:

Theorem 6. *Both sequences* $\left\{\alpha_k^{(\nu,\gamma)}\right\}$ *and* $\left\{c_n^{(\nu,\mu)}\right\}$ *are Hausdorff moment sequences for* $d = 1$ *and* $d = 2$.

3.3. Mean and Variance

The first and second order moments of these distributions can be easily computed as a consequence of the fact that they are compound distributions.

Theorem 7. *The distribution (10) of K verifies*

$$\begin{aligned}\sum_{k=0}^{+\infty}\alpha_k^{(\nu,\gamma)} &= 1,\\ EK &= \frac{d\gamma^2}{2\nu-2},\ \nu>1,\\ \sigma_K^2 &= \frac{d\gamma^2}{2(\nu-1)}\left(1+\frac{\gamma^2}{2}\frac{d+2(\nu-1)}{(\nu-1)(\nu-2)}\right),\ \nu>2.\end{aligned}$$

In the case of the sum of Student t- variables, we have

Theorem 8. *The distribution (11) of N verifies*

$$\begin{aligned}\sum_{k=0}^{+\infty}c_n^{(\nu,\mu)} &= 1,\\ EN &= \frac{d}{2}\left(\frac{B(\nu-1,\mu-1)}{B(\nu,\mu)}-1\right)\\ \sigma_N^2 &= \frac{d}{4}\left((d+2)\frac{B(\nu-2,\mu-2)}{B(\nu,\mu)}-2\frac{B(\nu-1,\mu-1)}{B(\nu,\mu)}-d\frac{B^2(\nu-1,\mu-1)}{B^2(\nu,\mu)}\right).\end{aligned}$$

3.4. Asymptotic behaviors

3.4.1. fixed value of γ

Let us now study the asymptotic behavior of the distribution (3) for large value of the degrees of freedom ν : note that the two next theorems can be checked directly by looking at the stochastic representation of K as given in Proposition 4, but their proofs are instructive by themselves.

Theorem 9. *For a fixed value of the parameter γ,*

$$\lim_{\nu\to+\infty}\alpha_k^{(\nu,\gamma)}=\begin{cases}1 & k=0\\ 0 & k\neq 0\end{cases}$$

We note that this result is a consequence of the fact that since $\mathbf{T}$ has covariance matrix $\frac{1}{2\nu-2}I_d$, the random variable $\mathbf{Z}=\mathbf{X}+\mathbf{T}$ has the same distribution as $\mathbf{X}$ as $\nu\to+\infty$.

3.4.2. covariance matrix of $\mathbf{T}$ fixed

Let us now look at the asymptotic behavior of these coefficients for large values of ν but for a fixed value of the covariance matrix of $\mathbf{T}$: since $\mathbf{T}$ has covariance matrix $\frac{1}{2(\nu-1)}I_d$, let us consider

$$\mathbf{Z}=\mathbf{X}+\eta\sqrt{2(\nu-1)}\mathbf{T}, \tag{12}$$

where $\mathbf{X}$ has covariance matrix $\sigma^2 I_d$ and $\eta\sqrt{2(\nu-1)}\mathbf{T}$ has covariance matrix $\eta^2 I_d$. Define

$$\tilde{\gamma}=\frac{\eta\sqrt{\nu-1}}{\sigma}$$

so that

$$f_{\mathbf{Z}}(\mathbf{z})=\sum_{k=0}^{+\infty}\alpha_k^{(\nu,\tilde{\gamma})}g_{k,\sigma}(\mathbf{z}). \tag{13}$$

Theorem 10. *The coefficients* $\alpha_k^{(\nu,\tilde{\gamma})}$ *in (13) behave for large* ν *as*

$$\alpha_k^{(\nu,\tilde{\gamma})} \sim \frac{\Gamma\left(k+\frac{d}{2}\right)}{k!\Gamma\left(\frac{d}{2}\right)} \left(\frac{\eta^2}{\sigma^2}\right)^k \left(1+\frac{\eta^2}{\sigma^2}\right)^{-k-\frac{d}{2}}$$

As a consequence, the density of $\mathbf{Z} = \mathbf{X} + \eta\sqrt{2(\nu-1)}\mathbf{T}$ *converges to a Gaussian density with covariance matrix* $\left(\sigma^2+\eta^2\right) I_d$*, as expected.*

In Fig.1 are shown the succesive terms of the approximation of the density of Z in the one-dimensional case ($d=1$): the upper curve is the density f_Z, the second one is the approximation of f_Z by the sum of the four first terms $0 \leq k \leq 3$ in (3), and the four lower curves are the successive terms of this sum for $k = 0, 1, 2$ and 3. The first ($k=0$) term (Gaussian density) fits f_Z around $z=0$, but the tail is underestimated; the other terms contribute to correct this underestimation, each adding, as k increases, a smaller contribution farther from the origin.



Figure 1: from top to bottom, density f_Z in the case $d=1$, its approximation by the sum of the first four term in (3) and these first four terms

4. Proofs

4.1. Proof of Theorem 1

Using a result by Keilson and Steutel [12], a stochastic representation for the random variable $\mathbf{Z}$ is

$$\mathbf{Z} = \sigma\mathbf{N}_1 + \frac{1}{\sqrt{2a}}\mathbf{N}_2 \overset{d}{=} \sqrt{\sigma^2+\frac{1}{2a}}\mathbf{N}$$

where $\mathbf{N}$ is $d-$variate standard normal, a is Gamma distributed $\Gamma(\nu,1)$ and $\overset{d}{=}$ denotes equality in distribution. The density of $\mathbf{Z}$ reads thus

$$f_{\mathbf{Z}}(\mathbf{z}) = \int_0^{+\infty} \frac{1}{\Gamma(\nu)} e^{-a} a^{\nu-1} \frac{1}{\left(2\pi\left(\sigma^2+\frac{1}{2a}\right)\right)^{\frac{d}{2}}} \exp\left(-\frac{\|\mathbf{z}\|^2}{2\left(\sigma^2+\frac{1}{2a}\right)}\right) da. \tag{14}$$

Since $\exp\left(-\frac{\|\mathbf{z}\|^2}{2\left(\sigma^2+\frac{1}{2a}\right)}\right) = \exp\left(-\frac{\|\mathbf{z}\|^2}{2\sigma^2}\right)\exp\left(\frac{\|\mathbf{z}\|^2}{2\sigma^2}\frac{\gamma^2}{a+\gamma^2}\right)$ with $\gamma^2 = \frac{1}{2\sigma^2}$, we deduce

$$\begin{aligned} f_{\mathbf{Z}}(\mathbf{z}) &= \frac{1}{\left(\sigma\sqrt{2\pi}\right)^d} \exp\left(-\frac{\|\mathbf{z}\|^2}{2\sigma^2}\right) \frac{1}{\Gamma(\nu)} \int_0^{+\infty} e^{-a} a^{\nu-1} \frac{a^{\frac{d}{2}}}{\left(a+\gamma^2\right)^{\frac{d}{2}}} \exp\left(\frac{\|\mathbf{z}^2\|}{2\sigma^2}\frac{\gamma^2}{a+\gamma^2}\right) da \\ &= \frac{1}{\left(\sigma\sqrt{2\pi}\right)^d} \exp\left(-\frac{\|\mathbf{z}\|^2}{2\sigma^2}\right) \frac{1}{\Gamma(\nu)} \sum_{k=0}^{+\infty} \frac{\gamma^{2k}}{k!} \left(\frac{\|\mathbf{z}\|^2}{2\sigma^2}\right)^k \int_0^{+\infty} e^{-a} a^{\nu+\frac{d}{2}-1} \left(a+\gamma^2\right)^{-k-\frac{d}{2}} da, \end{aligned}$$

where the interchange of the sum and the integral is justified by the Lebesgue monotone convergence theorem. Since, using [10, 4.642], $\forall k \in \mathbb{N}$,

$$\int_{\mathbb{R}^d} \left(\frac{\|\mathbf{x}\|^2}{2\sigma^2}\right)^k \exp\left(-\frac{\|\mathbf{x}\|^2}{2\sigma^2}\right) d\mathbf{x} = \frac{2\pi^{\frac{d}{2}}}{\Gamma\left(\frac{d}{2}\right)} \int_0^{+\infty} \left(\frac{r^2}{2\sigma^2}\right)^k \exp\left(-\frac{r^2}{2\sigma^2}\right) r^{d-1} dr,$$

which is equal to $\frac{\pi^{\frac{d}{2}}}{\Gamma(\frac{d}{2})}\left(\sigma\sqrt{2}\right)^d \Gamma\left(k+\frac{d}{2}\right)$, the function $g_{k,\sigma}$ as defined by (1) is a probability density and the result follows. Using the Kummer function

$$\Psi(\alpha,\beta;z) = \frac{1}{\Gamma(\alpha)} \int_0^{\infty} e^{-zt} t^{\alpha-1} (1+t)^{\beta-\alpha-1} dt, \tag{15}$$

cf. [10, p.1085], the integral in the expression of $f_{\mathbf{Z}}$ can be written

$$\int_0^{\infty} a^{\nu+\frac{d}{2}-1} e^{-a} (a+\gamma^2)^{-k-\frac{d}{2}} da = (\gamma^2)^{\nu-k}\Gamma(\nu+\frac{d}{2})\Psi(\nu+\frac{d}{2},\nu-k+1;\gamma^2). \tag{16}$$

4.2. Proof of Proposition 2

We have

$$\tilde{\mathbf{Z}} = a_2\left(\frac{a_1}{a_2}\mathbf{X}+\mathbf{T}\right) = a_2\left(\tilde{\sigma}\mathbf{N}_1+\mathbf{T}\right) = a_2\tilde{\mathbf{Z}}$$

with $\tilde{\sigma} = \frac{a_1}{a_2}\sigma$ so that $\tilde{\gamma} = \frac{a_2}{\sigma\sqrt{2}a_1}$ and

$$f_{\tilde{\mathbf{Z}}}(z) = \frac{1}{a_2^d} f_{\mathbf{Z}}\left(\frac{\mathbf{z}}{a_2}\right) = \sum_{k=0}^{+\infty} \alpha_k^{(\nu,\tilde{\gamma})} \frac{1}{a_2^d} g_{k,\frac{a_1}{a_2}\sigma}\left(\frac{\mathbf{z}}{a_2}\right)$$

but $\frac{1}{a_2^d} g_{k,\frac{a_1}{a_2}\sigma}\left(\frac{\mathbf{z}}{a_2}\right) = g_{k,a_1\sigma}(\mathbf{z})$ so that the result holds.

4.3. Proof of Theorem 3

Let us define $t = \frac{\sigma^2}{2}$ so that $\gamma = \frac{1}{\sigma\sqrt{2}} = \frac{1}{2\sqrt{t}}$ and consider the $k-$th term in the convex combination (3)

$$\alpha_k^{\left(\nu,\frac{1}{2\sqrt{t}}\right)} g_{k,\sqrt{2t}}(\mathbf{x}) = \frac{1}{\pi^{\frac{d}{2}}\Gamma(\nu)\,k!}\left(\frac{\|\mathbf{x}\|^2}{4t}\right)^k \exp\left(-\frac{\|\mathbf{x}\|^2}{4t}\right)\int_0^{+\infty}\frac{s^{\nu+\frac{d}{2}-1}\exp(-s)}{(1+4ts)^{k+\frac{d}{2}}}ds;$$

putting $w = 4ts$ and subordinating with the inverse Gamma distribution yields

$$\begin{aligned}\int_0^{+\infty}\alpha_k^{\left(\nu,\frac{1}{2\sqrt{t}}\right)} g_{k,\sqrt{2t}}(\mathbf{x})\,d\tilde{\gamma}_\mu(t) &= \frac{2^{-2\mu}}{\pi^{\frac{d}{2}}\Gamma(\mu)\Gamma(\nu)}\frac{\|\mathbf{x}\|^{2k}}{k!}\int_0^{+\infty}\left(\frac{1}{4t}\right)^{k+\nu+\frac{d}{2}}\frac{e^{-\frac{\|\mathbf{x}\|^2+1}{4t}}}{t^{\mu+1}}\int_0^{+\infty}\frac{w^{\nu+\frac{d}{2}-1}e^{-\frac{w}{4t}}}{(1+w)^{k+\frac{d}{2}}}dwdt\\ &= \frac{4^{-\mu-\nu-k-\frac{d}{2}}}{\pi^{\frac{d}{2}}\Gamma(\mu)\Gamma(\nu)}\frac{\|\mathbf{x}\|^{2k}}{k!}\int_0^{+\infty}\frac{w^{\nu+\frac{d}{2}-1}}{(1+w)^{k+\frac{d}{2}}}\int_0^{+\infty}\frac{e^{-\frac{w+\|\mathbf{x}\|^2+1}{4t}}}{t^{k+\mu+\nu+\frac{d}{2}+1}}dtdw.\end{aligned}$$

By the change of variable $u = \frac{w+\|\mathbf{x}\|^2+1}{4t}$, this is equal to

$$\frac{1}{4^{\mu+\nu+k+\frac{d}{2}}\pi^{\frac{d}{2}}\Gamma(\mu)\Gamma(\nu)}\frac{\|\mathbf{x}\|^{2k}}{k!}\int_0^{+\infty}\frac{w^{\nu+\frac{d}{2}-1}}{(1+w)^{k+\frac{d}{2}}}\left(\frac{w+\|\mathbf{x}\|^2+1}{4}\right)^{-k-\mu-\nu-\frac{d}{2}}\int_0^{+\infty}u^{k+\mu+\nu+\frac{d}{2}-1}e^{-u}dudw$$

and, since the latter integral is a Gamma integral, to

$$\frac{\Gamma\left(\mu+\nu+k+\frac{d}{2}\right)}{\pi^{\frac{d}{2}}\Gamma(\mu)\Gamma(\nu)}\frac{\|\mathbf{x}\|^{2k}}{k!}\int_0^{+\infty}\frac{w^{\nu+\frac{d}{2}-1}}{(1+w)^{k+\frac{d}{2}}(w+\|\mathbf{x}\|^2+1)^{k+\mu+\nu+\frac{d}{2}}}dw$$

The latter integral can be identified as Euler's integral representation of a hypergeometric function [3, Th. 2.2.1 p.65]

$$\begin{aligned}&\int_0^{+\infty} \frac{w^{\nu+\frac{d}{2}-1}}{(1+w)^{k+\frac{d}{2}}(w+\|\mathbf{x}\|^2+1)^{k+\mu+\nu+\frac{d}{2}}}dw \\ =\ & \frac{\Gamma\left(2k+\mu+\frac{d}{2}\right)\Gamma\left(\nu+\frac{d}{2}\right)}{\Gamma(2k+\mu+\nu+d)} {}_2F_1\left(\begin{array}{c} k+\mu+\nu+\frac{d}{2},\ 2k+\mu+\frac{d}{2} \\ 2k+\mu+\nu+d \end{array};-\|\mathbf{x}\|^2\right)\end{aligned}$$

so that, using Pfaff's formula [3, 2.2.6 p.68]

$${}_2F_1\left(\begin{array}{c} a,\ b \\ c \end{array};x\right) = (1-x)^{-a}\,{}_2F_1\left(\begin{array}{c} a,\ c-b \\ c \end{array};\frac{x}{x-1}\right)$$

we deduce

$$\begin{aligned}\int_0^{+\infty}\alpha_k^{\left(\nu,\frac{1}{2\sqrt{t}}\right)} g_{k,\sqrt{2t}}(\mathbf{x})\,d\tilde{\gamma}_\mu(t) &= \frac{\Gamma\left(\mu+\nu+k+\frac{d}{2}\right)\Gamma\left(\nu+\frac{d}{2}\right)}{\pi^{\frac{d}{2}}\Gamma(\mu)\Gamma(\nu)\,k!}\frac{\Gamma\left(2k+\mu+\frac{d}{2}\right)}{\Gamma(2k+\mu+\nu+d)} \\ &\times \frac{\|\mathbf{x}\|^{2k}}{(1+\|\mathbf{x}\|^2)^{k+\mu+\nu+\frac{d}{2}}}\,{}_2F_1\left(\begin{array}{c} k+\mu+\nu+\frac{d}{2},\ \nu+\frac{d}{2} \\ 2k+\mu+\nu+d \end{array};\frac{\|\mathbf{x}\|^2}{1+\|\mathbf{x}\|^2}\right).\end{aligned}$$

Replacing the hypergeometric function by its series, we deduce

$$\begin{aligned}\int_0^{+\infty}\alpha_k^{\left(\nu,\frac{1}{2\sqrt{t}}\right)} g_{k,\sqrt{2t}}(\mathbf{x})\,d\tilde{\gamma}_\mu(t) &= \frac{\Gamma\left(\mu+\nu+k+\frac{d}{2}\right)\Gamma\left(\nu+\frac{d}{2}\right)}{\pi^{\frac{d}{2}}\Gamma(\mu)\Gamma(\nu)\,k!}\frac{\Gamma\left(2k+\mu+\frac{d}{2}\right)}{\Gamma(2k+\mu+\nu+d)} \\ &\times \sum_{l=0}^{+\infty}\frac{\left(\nu+\frac{d}{2}\right)_l\left(k+\mu+\nu+\frac{d}{2}\right)_l}{(2k+\mu+\nu+d)_l\,l!}\frac{\|\mathbf{x}\|^{2k+2l}}{(1+\|\mathbf{x}\|^2)^{k+l+\mu+\nu+\frac{d}{2}}} \\ &= \sum_{l=0}^{+\infty} d_{k,l}\phi_{l+k,\eta}(\mathbf{x})\end{aligned}$$

where $\eta=\mu+\nu$, the density $\phi_{n,\eta}(\mathbf{x})$ is defined by (2) and the coefficient $d_{k,l}$ is equal to

$$\begin{aligned}d_{k,l} &= \frac{\Gamma\left(\mu+\nu+k+\frac{d}{2}\right)\Gamma\left(\nu+\frac{d}{2}\right)}{\pi^{\frac{d}{2}}\Gamma(\mu)\Gamma(\nu)\,k!}\frac{\Gamma\left(2k+\mu+\frac{d}{2}\right)}{\Gamma(2k+\mu+\nu+d)}\frac{\left(\nu+\frac{d}{2}\right)_l\left(k+\mu+\nu+\frac{d}{2}\right)_l}{(2k+\mu+\nu+d)_l\,l!}\frac{\Gamma(\mu+\nu)\Gamma\left(k+l+\frac{d}{2}\right)}{\Gamma\left(\mu+\nu+k+l+\frac{d}{2}\right)}\frac{\pi^{\frac{d}{2}}}{\Gamma\left(\frac{d}{2}\right)} \\ &= \frac{\Gamma\left(\nu+\frac{d}{2}\right)\left(\nu+\frac{d}{2}\right)_l}{\Gamma(\mu)\Gamma(\nu)\,k!l!}\frac{\Gamma\left(2k+\mu+\frac{d}{2}\right)\Gamma(\mu+\nu)\Gamma\left(k+l+\frac{d}{2}\right)}{\Gamma(2k+l+\mu+\nu+d)\Gamma\left(\frac{d}{2}\right)}.\end{aligned}$$

using the identity $(a)_l\Gamma(a)=\Gamma(a+l)$ twice. Writing $\Gamma\left(2k+\mu+\frac{d}{2}\right)=\left(\mu+\frac{d}{2}\right)_{2k}\Gamma\left(\mu+\frac{d}{2}\right)$, $\Gamma\left(k+l+\frac{d}{2}\right)=\left(\frac{d}{2}\right)_{k+l}\Gamma\left(\frac{d}{2}\right)$ and $\Gamma(2k+l+\mu+\nu+d)=(\mu+\nu+d)_{2k+l}\Gamma(\mu+\nu+d)$, we deduce

$$d_{k,l}=\frac{B\left(\nu+\frac{d}{2},\mu+\frac{d}{2}\right)}{B(\nu,\mu)\,k!l!}\frac{\left(\mu+\frac{d}{2}\right)_{2k}\left(\nu+\frac{d}{2}\right)_l\left(\frac{d}{2}\right)_{k+l}}{(\mu+\nu+d)_{2k+l}}.$$

Now the density of $\mathbf{Y}=\mathbf{T}_1+\mathbf{T}_2$ is the sum over k of these terms and reads

$$f_Y(\mathbf{x})=\sum_{k,l=0}^{+\infty} d_{k,l}\phi_{k+l,\eta}(\mathbf{x}).$$

We perform the change of variable $n=k+l$ to obtain

$$f_{\mathbf{Y}}(\mathbf{x})=\sum_{n=0}^{+\infty}\phi_{n,\eta}(\mathbf{x})\sum_{k=0}^{n} d_{k,n-k}.$$

The inner sum can be written

$$\begin{aligned} c_n^{(\nu,\mu)} = \sum_{k=0}^{n} d_{k,n-k} &= \frac{B\left(\nu+\frac{d}{2},\mu+\frac{d}{2}\right)}{B\left(\nu,\mu\right)} \frac{\left(\frac{d}{2}\right)_n}{n!} \sum_{k=0}^{n} \binom{n}{k} \frac{\left(\mu+\frac{d}{2}\right)_{2k} \left(\nu+\frac{d}{2}\right)_{n-k}}{\left(\mu+\nu+d\right)_{n+k}} \\ = \frac{B\left(\nu+\frac{d}{2},\mu+\frac{d}{2}\right)}{B\left(\nu,\mu\right)} & \frac{\left(\frac{d}{2}\right)_n}{\left(\mu+\nu+d\right)_{2n} n!} \sum_{k=0}^{n} \binom{n}{k} \left(\mu+\frac{d}{2}\right)_{2k} \left(\nu+\frac{d}{2}\right)_{n-k} \left(\mu+\nu+d+n+k\right)_{n-k} \end{aligned}$$

where we have used the identity

$$\frac{1}{\left(\mu+\nu+d\right)_{n+k}} = \frac{\left(\mu+\nu+d+n+k\right)_{n-k}}{\left(\mu+\nu+d\right)_{2n}}.$$

By the Chu-Vandermonde identity [3, p.70]

$$\begin{aligned} \left(\mu+\nu+d+n+k\right)_{n-k} &= \left(\left(\mu+\frac{d}{2}+2k\right)+\left(\nu+\frac{d}{2}+n-k\right)\right)_{n-k} \\ &= \sum_{j=0}^{n-k} \binom{n-k}{j} \left(\mu+\frac{d}{2}+2k\right)_{n-k-j} \left(\nu+\frac{d}{2}+n-k\right)_{j} \end{aligned}$$

and using the identities $\left(\mu+\frac{d}{2}\right)_{2k}\left(\mu+\frac{d}{2}+2k\right)_{n-k-j} = \left(\mu+\frac{d}{2}\right)_{n+k-j}$ and $\left(\nu+\frac{d}{2}\right)_{n-k}\left(\nu+\frac{d}{2}+n-k\right)_{j} = \left(\nu+\frac{d}{2}\right)_{n-k+j}$, we deduce

$$\begin{aligned} c_n^{(\nu,\mu)} &= \frac{B\left(\nu+\frac{d}{2},\mu+\frac{d}{2}\right)}{B\left(\nu,\mu\right)} \frac{\left(\frac{d}{2}\right)_n}{\left(\mu+\nu+d\right)_{2n} n!} \sum_{k=0}^{n} \sum_{j=0}^{n-k} \binom{n}{k} \binom{n-k}{j} \left(\mu+\frac{d}{2}\right)_{n+k-j} \left(\nu+\frac{d}{2}\right)_{n-k+j} \\ &= \frac{B\left(\nu+\frac{d}{2},\mu+\frac{d}{2}\right)}{B\left(\nu,\mu\right)} \frac{\left(\frac{d}{2}\right)_n}{n!} \sum_{p=-n}^{n} \binom{n}{p}_2 \frac{\left(\mu+\frac{d}{2}\right)_{n+p} \left(\nu+\frac{d}{2}\right)_{n-p}}{\left(\mu+\nu+d\right)_{2n}}, \end{aligned}$$

where we have used the change of summation index $p=k-j$ and where the trinomial [2] coefficient $\binom{n}{p}_2$ is

$$\binom{n}{p}_2 = \sum_{\substack{j,k=0 \\ j+k\le n;\, k-j=p}}^{n} \binom{n}{k} \binom{n-k}{j}, \ -n \le p \le n.$$

Writing $\left(\mu+\frac{d}{2}\right)_{n+p} \Gamma\left(\mu+\frac{d}{2}\right) = \Gamma\left(\mu+n+p+\frac{d}{2}\right)$ and $\left(\nu+\frac{d}{2}\right)_{n-p} \Gamma\left(\nu+\frac{d}{2}\right) = \Gamma\left(\nu+\frac{d}{2}+n-p\right)$ and replacing the resulting Beta function by its integral representation we obtain

$$c_n^{(\nu,\mu)} = \frac{1}{B\left(\nu,\mu\right)} \frac{\left(\frac{d}{2}\right)_n}{n!} \sum_{p=-n}^{n} \binom{n}{p}_2 \int_0^1 t^{\mu+n+p+\frac{d}{2}-1} \left(1-t\right)^{\nu+n-p+\frac{d}{2}-1} dt,$$

and since the generating function of the trinomial coefficient is

$$\sum_{p=-n}^{+n} \binom{n}{p}_2 x^p = \left(x+x^{-1}+1\right)^n$$

we finally obtain

$$c_n^{(\nu,\mu)} = \frac{1}{B\left(\nu,\mu\right)} \frac{\left(\frac{d}{2}\right)_n}{n!} \int_0^1 t^{\mu+n+\frac{d}{2}-1} \left(1-t\right)^{\nu+n+\frac{d}{2}-1} \left(1+\frac{t}{1-t}+\frac{1-t}{t}\right)^n dt$$

which is the desired result.

4.4. Proof of Theorem 6

In the case of the coefficients $\alpha_k^{(\nu,\gamma)}$, we can factorize

$$\alpha_k^{(\nu,\gamma)} = \frac{\left(\frac{d}{2}\right)_k}{k!}\tau_k$$

where

$$\tau_k = \frac{\left(\sigma\sqrt{2}\right)^d}{\Gamma(\nu)}\int_0^{+\infty}\frac{s^{\nu+\frac{d}{2}-1}e^{-s}}{\left(1+2\sigma^2 s\right)^{k+\frac{d}{2}}}ds = \frac{1}{\left(2\sigma^2\right)^{\nu}\Gamma(\nu)}\int_0^1 u^k\frac{(1-u)^{\nu+\frac{d}{2}-1}}{u^{\nu+1}}\exp\left(-\frac{1-u}{2\sigma^2 u}\right)du.$$

This expression shows that the sequence $\{\tau_k\}$ is also a Hausdorff moment sequence (not normalized) and hence $\alpha_k^{(\nu,\gamma)}$ is a Hausdorff moment sequence for $d=1,2$ as product of two such sequences. It is not normalized since $\sum_{k=0}^{+\infty}\alpha_k^{(\nu,\gamma)} = 1$. In fact, $\frac{\left(\frac{d}{2}\right)_k}{k!}$ is trivially a Hausdorff moment sequence when $d=2$, and for $d=1$ we have

$$\frac{\left(\frac{1}{2}\right)_k}{k!} = \frac{1}{\pi}\int_0^1\frac{s^k ds}{\sqrt{s(1-s)}}.$$

It is a general fact that $s_k = \frac{(a)_k}{(b)_k}$ is a Hausdorff moment sequence when $0 < a \le b$, see [5, formula (26)] and if $a > b$ then s_k is not even a Hamburger moment sequence since the Hankel determinant $D_2 = s_0 s_2 - s_1^2 = \frac{a(b-a)}{b^2(b+1)} < 0$.

This shows that the density (3) is an infinite convex combination of the densities (1) and the discrete probability $\alpha^{(\nu,\gamma)} = \sum_{k=0}^{+\infty}\alpha_k^{(\nu,\gamma)}\delta_k$ is infinitely divisible for $d=1,2$ since the sequence $\left\{\alpha_k^{(\nu,\gamma)}\right\}$ is completely monotonic, which is the same as being a Hausdorff moment sequence.

In the case of the sequence $\left\{c_n^{(\nu,\mu)}\right\}$, we find

$$\begin{aligned} c_n^{(\nu,\mu)} &= \frac{1}{B(\nu,\mu)}\frac{\left(\frac{d}{2}\right)_n}{n!}\int_{\frac{3}{4}}^1 u^n \times \\ &\left[\left(\frac{1}{2}-\sqrt{u-\frac{3}{4}}\right)^{\mu+\frac{d}{2}-1}\left(\frac{1}{2}+\sqrt{u-\frac{3}{4}}\right)^{\nu+\frac{d}{2}-1} + \left(\frac{1}{2}+\sqrt{u-\frac{3}{4}}\right)^{\mu+\frac{d}{2}-1}\left(\frac{1}{2}-\sqrt{u-\frac{3}{4}}\right)^{\nu+\frac{d}{2}-1}\right]\frac{du}{2\sqrt{u-\frac{3}{4}}}. \end{aligned}$$

As before $\left\{c_n^{(\nu,\mu)}\right\}$ is a Hausdorff moment sequence for $d=1,2$.

4.5. Proof of Theorem 7

Although the first result can be obtained by integrating (3) over $\mathbf{z}\in\mathbb{R}^d$, we check it directly since the computation of the sum is instructive by itself. For any $u \in \left]-1,1\right[$, we have by the binomial series

$$\sum_{k=0}^{+\infty}\frac{\Gamma\left(k+\frac{d}{2}\right)}{k!}u^k = \frac{\Gamma\left(\frac{d}{2}\right)}{(1-u)^{\frac{d}{2}}} \tag{17}$$

so that, with $u = \frac{\gamma^2}{a+\gamma^2}$,

$$\sum_{k=0}^{+\infty}\frac{\Gamma\left(k+\frac{d}{2}\right)}{k!}\left(\frac{\gamma^2}{a+\gamma^2}\right)^k = \Gamma\left(\frac{d}{2}\right)\left(\frac{a+\gamma^2}{a}\right)^{\frac{d}{2}}$$

and we deduce

$$\sum_{k=0}^{+\infty}\alpha_k^{(\nu,\gamma)} = \int_0^{+\infty}\frac{e^{-a}}{\Gamma(\nu)\Gamma\left(\frac{d}{2}\right)}\frac{a^{\nu+\frac{d}{2}-1}}{\left(a+\gamma^2\right)^{\frac{d}{2}}}\Gamma\left(\frac{d}{2}\right)\left(\frac{a+\gamma^2}{a}\right)^{\frac{d}{2}}da = 1.$$

The expectation and variances are easily computed from the stochastic representation of $K = NB\left(\frac{d}{2}, \frac{a}{1+a}\right)$; the mean of a negative binomial variable $NB(r,p)$ being $\frac{r(1-p)}{p}$,

$$EK = \frac{d}{2}E\frac{\frac{1}{1+a}}{\frac{a}{1+a}} = \frac{d}{2}E\frac{1}{a} = \frac{d\gamma^2}{2\nu-2}$$

since a is $\Gamma\left(\nu,\gamma^2\right)$. Since moreover, the second order moment of a negative binomial is $\frac{r(1-p)(1+r(1-p))}{p^2}$, a straightforward computation gives

$$\begin{aligned} EK^2 &= \frac{d}{2}\left(E\frac{1+a}{a^2} + \frac{d}{2}E\frac{1}{a^2}\right) = \frac{d}{2}\left(1+\frac{d}{2}\right)E\frac{1}{a^2} + \frac{d}{2}E\frac{1}{a} \\ &= \frac{d}{2}\left(1+\frac{d}{2}\right)\frac{\gamma^4}{(\nu-1)(\nu-2)} + \frac{d}{2}\frac{\gamma^2}{\nu-1} \end{aligned}$$

and

$$\sigma_K^2 = \frac{d}{2}\frac{\gamma^2}{\nu-1}\left(1+\frac{\gamma^2}{2}\frac{d+2(\nu-1)}{(\nu-1)(\nu-2)}\right).$$

4.6. Proof of Theorem 8

The fact that the coefficients $c_n^{(\nu,\mu)}$ sum to 1 is proved in the same way as in the Proof of Theorem 7; the expectation is computed accordingly as

$$EN = \frac{d}{2}E\left(\frac{1}{t(1-t)} - 1\right)$$

with

$$E\frac{1}{t(1-t)} = \frac{B(\nu-1,\mu-1)}{B(\nu,\mu)}.$$

The second order moment is similarly computed as

$$EN^2 = \frac{d}{2}E\frac{(1-t(1-t))\left(1+\frac{d}{2}(1-t(1-t))\right)}{t^2(1-t)^2} = \frac{d}{2}\left(1+\frac{d}{2}\right)E\frac{1}{t^2(1-t)^2} - \frac{d(1+d)}{2}E\frac{1}{t(1-t)} + \frac{d^2}{4}$$

so that the variance reads

$$\sigma_N^2 = \frac{d^2}{4} - \frac{d(1+d)}{2}\frac{B(\nu-1,\mu-1)}{B(\nu,\mu)} + \frac{d}{2}\left(1+\frac{d}{2}\right)\frac{B(\nu-2,\mu-2)}{B(\nu,\mu)} - \frac{d^2}{4}\left(\frac{B(\nu-1,\mu-1)}{B(\nu,\mu)} - 1\right)^2$$

and the result follows after simplification.

4.7. Proof of Theorem 9

It is clearly enough to prove that $\lim_{\nu\to\infty}\alpha_0^{(\nu,\gamma)} = 1$ since $0 \le \alpha_k^{(\nu,\gamma)} \le 1-\alpha_0^{(\nu,\gamma)}$ for $k>0$. We have

$$\alpha_0^{(\nu,\gamma)} = \frac{1}{\Gamma(\nu)}\int_0^\infty e^{-a}a^{\nu-1}\left(1+\frac{\gamma^2}{a}\right)^{-d/2} da,$$

hence

$$1-\alpha_0^{(\nu,\gamma)} = \frac{1}{\Gamma(\nu)}\int_0^\infty e^{-a}a^{\nu-1}\left(1-\left(1+\frac{\gamma^2}{a}\right)^{-d/2}\right) da = \frac{d\gamma}{\Gamma(\nu)}\int_0^\infty e^{-a}a^{\nu-2}\xi\left(1+\frac{\xi^2}{a}\right)^{-d/2-1} da,$$

for some $0<\xi<\gamma$ by the mean value theorem. For $\nu>1$ we therefore get

$$1-\alpha_0^{(\nu,\gamma)} = \frac{d\gamma^2}{\Gamma(\nu)}\int_0^\infty e^{-a}a^{\nu-2}\, da = \frac{d\gamma^2}{\nu-1}$$

which tends to 0 for $\nu\to\infty$.

4.8. Proof of Theorem 10

The coefficients $\alpha_k^{(\nu,\tilde{\gamma})}$ read

$$\begin{aligned}
\alpha_k^{(\nu,\tilde{\gamma})} &= \frac{\Gamma(k+\frac{d}{2})}{k!\Gamma\left(\frac{d}{2}\right)\Gamma(\nu)}\left(\frac{\eta^2\left(\nu-1\right)}{\sigma^2}\right)^k\int_0^{+\infty}\exp(-a)a^{\nu+\frac{d}{2}-1}\left(a+\frac{\eta^2\left(\nu-1\right)}{\sigma^2}\right)^{-k-\frac{d}{2}}da \\
&= \frac{\Gamma(k+\frac{d}{2})}{k!\Gamma\left(\frac{d}{2}\right)\Gamma(\nu)}\left(\frac{\eta^2}{\sigma^2}\right)^k\left(\nu-1\right)^{\nu}\int_0^{+\infty}\exp(-b\left(\nu-1\right))b^{\nu+\frac{d}{2}-1}\left(b+\frac{\eta^2}{\sigma^2}\right)^{-k-\frac{d}{2}}db
\end{aligned}$$

using change of variable $a=b\left(\nu-1\right)$. Let us apply Laplace's method [16, p.278]: if $f'\left(x_0\right)=0$ and $f''\left(x_0\right)<0$ then for large M,

$$\int_a^b g(x)e^{Mf(x)}dx \sim g\left(x_0\right)e^{Mf\left(x_0\right)}\sqrt{\frac{2\pi}{M|f''\left(x_0\right)|}}.$$

Choosing $M=\nu-1$, $f\left(x\right)=-x+\log x$, $g(x)=x^{\frac{d}{2}}\left(x+a^2\right)^{-k-\frac{d}{2}}$ and $x_0=1$, an equivalent of the right-hand side integral reads

$$e^{-\left(\nu-1\right)}\sqrt{\frac{2\pi}{\nu-1}}\left(1+\frac{\eta^2}{\sigma^2}\right)^{-k-\frac{d}{2}}.$$

Thus an equivalent of the coefficients $\alpha_k^{(\nu,\tilde{\gamma})}$ reads

$$\frac{\Gamma(k+\frac{d}{2})}{k!\Gamma\left(\frac{d}{2}\right)}\frac{\left(\nu-1\right)^{\nu-\frac{1}{2}}e^{-\left(\nu-1\right)}\sqrt{2\pi}}{\Gamma\left(\nu\right)}\left(\frac{\eta^2}{\sigma^2}\right)^k\left(1+\frac{\eta^2}{\sigma^2}\right)^{-k-\frac{d}{2}}$$

which, by Stirling's formula, is equivalent to

$$\frac{\Gamma\left(k+\frac{d}{2}\right)}{k!\Gamma\left(\frac{d}{2}\right)}\left(\frac{\eta^2}{\sigma^2}\right)^k\left(1+\frac{\eta^2}{\sigma^2}\right)^{-k-\frac{d}{2}}.$$

As a consequence, for large values of ν, the density $f_{\mathbf{Z}}$ of $\mathbf{Z}$ reads

$$\begin{aligned}
f_{\mathbf{Z}}\left(\mathbf{z}\right) &\sim \sum_{k=0}^{+\infty}\frac{\Gamma\left(\frac{d}{2}\right)}{\left(\sigma\sqrt{2\pi}\right)^d\Gamma\left(k+\frac{d}{2}\right)}\left(\frac{\|\mathbf{z}\|^2}{2\sigma^2}\right)^k\exp\left(-\frac{\|\mathbf{z}\|^2}{2\sigma^2}\right)\frac{\Gamma\left(k+\frac{d}{2}\right)}{k!\Gamma\left(\frac{d}{2}\right)}\left(\frac{\eta^2}{\sigma^2}\right)^k\left(1+\frac{\eta^2}{\sigma^2}\right)^{-k-\frac{d}{2}} \\
&= \frac{1}{\left(\sigma\sqrt{2\pi}\right)^d}\frac{1}{\left(1+\frac{\eta^2}{\sigma^2}\right)^{\frac{d}{2}}}\exp\left(-\frac{\|\mathbf{z}\|^2}{2\sigma^2}\right)\exp\left(\frac{\|\mathbf{z}\|^2}{2\sigma^2}\frac{\frac{\eta^2}{\sigma^2}}{1+\frac{\eta^2}{\sigma^2}}\right) \\
&= \frac{1}{\left(\tilde{\sigma}\sqrt{2\pi}\right)^d}\exp\left(-\frac{\|\mathbf{z}\|^2}{2\tilde{\sigma}^2}\right)
\end{aligned}$$

with

$$\tilde{\sigma}^2=\sigma^2+\eta^2.$$

4.9. Proof of Proposition 4

Conditionnally to a given value of $K=k$, the random variable $\sigma\|\mathbf{X}_{2k+d}\|\mathbf{U}_d$ follows the distribution $g_{k,\sigma}$; let us check that the random variable K follows a $NB\left(\frac{d}{2},\frac{a}{1+a}\right)$ compound distribution where a is $\Gamma\left(\nu,\gamma^2\right)$: conditionnally to a, such a compound distribution reads

$$p_k|a=\binom{k+\frac{d}{2}-1}{k}\left(\frac{a}{1+a}\right)^{\frac{d}{2}}\left(\frac{1}{1+a}\right)^k$$

so that

$$\alpha_k^{(\nu,\gamma)} = Ep_k|a = \binom{k+\frac{d}{2}-1}{k} \int_0^{+\infty} \frac{\gamma^{2\nu}}{\Gamma(\nu)} e^{-\gamma^2 a} a^{\nu+\frac{d}{2}-1} (1+a)^{-k-\frac{d}{2}} da$$
$$= \frac{\Gamma\left(k+\frac{d}{2}\right)\gamma^{2k}}{k!\Gamma\left(\frac{d}{2}\right)\Gamma(\nu)} \int_0^{+\infty} e^{-b} b^{\nu+\frac{d}{2}-1} \left(\gamma^2+b\right)^{-k-\frac{d}{2}} db$$

by the change of variable $b = \gamma^2 a$; this expression coincides with (4).

4.10. Proof of Proposition 5

Conditionnally to a value of $N = n$, the random variable $\|\mathbf{T}_{2n+d}\|\mathbf{U}_d$ follows the distribution $\phi_{n,\nu+\mu}$; we check that the random variable N follows a $NB\left(\frac{d}{2}, t(1-t)\right)$, where t follows a Beta distribution with parameters μ and ν: conditionnally to t, this compound distribution reads

$$p_n|t = \binom{n+\frac{d}{2}-1}{n} t^{\frac{d}{2}} (1-t)^{\frac{d}{2}} (1-t(1-t))^n$$

so that

$$c_n^{(\nu,\mu)} = Ep_n|t = \frac{1}{B(\nu,\mu)} \binom{n+\frac{d}{2}-1}{n} \int_0^1 t^{\mu+\frac{d}{2}-1} (1-t)^{\nu+\frac{d}{2}-1} (1-t(1-t))^n dt$$
$$= \frac{1}{B(\nu,\mu)} \frac{\Gamma\left(n+\frac{d}{2}\right)}{n!\Gamma\left(\frac{d}{2}\right)} \int_0^1 t^{\mu+\frac{d}{2}-1} (1-t)^{\nu+\frac{d}{2}-1} (1-t(1-t))^n dt$$

which coincides with (7).

Acknowledgements

The work of the first author has been supported by grant 272-07-0321 from the Danish Research Council for Nature and Universe. CV thanks CB for his invitation to the University of Copenhagen in February 2009. The paper was finished during a stay of the first author as Guest Professor at University of Marne la Vallée in May 2009.

References

[1] M. Abramowitz, I. Stegun, Handbook of Mathematical Functions: with formulas, Graphs, and Mathematical Tables, Dover Publications, 1965.

[2] G. Andrews, Euler's 'exemplum memorabile inductionis fallacis' and $q-$Trinomial Coefficients, J. Amer. Math. Soc. 3, 653–669, 1990.

[3] G. E. Andrews, R. Askey, and R. Roy, Special Functions, Cambridge University Press, Cambridge, UK, 1st ed., Feb. 1999.

[4] V. E. Bening, V. Yu. Korolev, On an application of the Student distribution in the theory of probability and mathematical statistics, Theory Probab. Appl., Vol. 49, No. 3 (2005), 377–391.

[5] C. Berg, On powers of Stieltjes moment sequences, II, J. Comput. Appl. Math., Vol. 199 (2007), 23–38.

[6] C. Berg, C. Vignat, Linearization Coefficients of Bessel Polynomials and Properties of Student t-Distributions, Constructive Approximation, Vol. 27 (2008), 15–32.

[7] C. Berg, C. Vignat, On some results of Cufaro Petroni about Student t-processes, J. Phys. A: Math. Theor. 41 (2008) 265004.

[8] N. Cufaro Petroni, Mixtures in nonstable Lévy processes, J. Phys. A: Math. Theor. 40 (2007), 2227–2250.

[9] K.-T. Fang, S. Kotz, K. W. Ng, Symmetric Multivariate and Related Distributions, Monographs on Statistics and Applied Probability, Chapman & Hall/CRC, 1989.

[10] I. S. Gradshteyn, I. M. Ryzhik, Table of Integrals, Series, and Products, Seventh Edition, Academic Press, 2007.

[11] C. C. Heyde, N. N. Leonenko, Student Processes, Adv. Appl. Prob. 37 (2005), 342–365.

[12] J. Keilson, F.W. Steutel, Mixtures of Distributions, Moment Inequalities and Measures of Exponentiality and Normality, Annals of Probability, Volume 2, Issue 1, Feb. 1974, 112–130.

[13] G. P. Nason, On the sum of t and Gaussian random variables, Statistics and Probability Letters 76 (2006) 1280.

[14] H. Ruben, On the distribution of the weighted difference of two independent Student variables. J. Roy. Statist. Soc. Ser. B, 22 (1960), 188–194.

[15] G.A. Walker, J.G. Saw, The Distribution of Linear Combinations of t-Variables. Journal of the American Statistical Association, Vol. 73, Issue 364, Dec. 1978, 876–878.

[16] D.V. Widder, The Laplace Transform, Princeton University Press, 1946.

[17] V. Witkovský, Exact distribution of positive linear combinations of inverted chi-square random variables with odd degrees of freedom. Statistics and Probability Letters 56 (2002), 45–50.